CORRESPONDENCE

Open Access

Check for

Is nitric oxide a clue to endemic goitre in highlanders?

Jon O. Lundberg^{1*} and Eddie Weitzberg¹

Abstract

Goitre is commonly caused by a lack of iodine in the diet. This condition is particularly prevalent in high-altitude areas where iodine deficiency is common. Here we speculate that inorganic nitrate, the oxidation product of nitric oxide, which is generated endogenously at very high levels in highlanders, further increases the risk of goitre and thyroid dysfunction in this population by inhibiting the transport of iodide into the thyroid gland. Indeed, it is well-known that nitrate and iodide compete for such transport. While iodine scarcity is a primary cause of goitre, the excessive nitrate levels in highlanders may further hinder iodide transport, exacerbating the problem.

Keywords Goitre, Nitrate, Nitric oxide, Highlanders, Thyroid, Iodide

Sir,

The causes of goitre are multifold but a lack of iodine (I_2) in the diet is common worldwide [1]. People living at high altitudes are at greater risk of goitre and prior to the controlled introduction of iodine supplementation through salt, oil and some other vehicles, the incidence in some Himalayan areas could approach 100% in the younger population [2]. Iodide (I⁻), the ionic form of iodine, is an essential element of thyroid hormones and is actively transported into the thyroid gland via a specific transporter, the sodium/iodide symporter, in a process also dependent on ATP, iron and magnesium. It is known that such transport occurs in competition with some other anions including thiocyanate (SCN⁻) and nitrate (NO_3^{-}) [3]. Diets rich in such compounds are sometimes referred to as goitrogenic, as they may lead to a decreased transport of iodide with resulting goitre and thyroid dysfunction [3, 4]. Nitrate is not only found in our diet, it is

also generated endogenously being an oxidation product of nitric oxide (NO) [5]. A report published some years ago showed that endogenous NO generation by endothelial NO synthase (eNOS) is exceptionally high in Tibetans [6]. In fact, plasma levels of nitrate in these highlanders are more than 10 times higher than in lowlanders and it was speculated that such massive NO production helps to offset high-altitude hypoxia and to achieve normal oxygen delivery through the vasodilatory effects of NO [6]. We speculate that the persistently high circulating levels of nitrate in highlanders may come with an unwanted side effect, namely inhibition of iodide transport and increased risk of goitre and thyroid dysfunction (Fig. 1). This may help to explain why historically, endemic goitre has classically been seen in high-altitude areas [2]. In addition, the constant exposure to hypoxia may also affect mitochondrial function which could indirectly affect iodide uptake through ATP-dependent processes. While a scarcity of dietary iodine remains the

*Correspondence:

Jon O. Lundberg

jon.lundberg@ki.se

¹ Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



Fig. 1 Historically, goitre has been common in highland areas where it could become endemic. This is mainly because the soil is poor in iodine but another mechanism may also be in play. Endogenous generation of nitric oxide (NO) is exceptionally high in highlanders due to continuous high activation of endothelial NO synthase (eNOS) in response to the low oxygen at these altitudes. In the body NO is rapidly converted to nitrate (NO_3^-), an anion known to compete with iodide (Γ) for uptake in the thyroid gland. Here we propose that the high susceptibility to goitre in highlanders may be partly related to the persistently high endogenous formation of NO and nitrate in these populations

NOS 1

major factor explaining endemic goitre, the additional mechanism discussed here may contribute, in particular in highlanders. In a situation with already little iodine coming from the diet, the endogenously-derived nitrate may become overwhelming, thereby further preventing transport of iodide into the thyroid gland for production of thyroid hormones.

Authors' contributions

Both authors conceived the study and together wrote the letter.

Funding

We have received grants from the Swedish Research Council (VR) and the Swedish Heart and Lung Foundation (HLF).

Availability of data and materials N/A.

Declarations

Ethics approval and consent to participate N/A.

Consent for publication

Both authors gave their consent to publish this article.

Competing interests

The authors declare no competing interests.

Received: 26 June 2023 Accepted: 24 September 2023 Published online: 02 October 2023

References

- Zimmermann MB, Boelaert K. lodine deficiency and thyroid disorders. Lancet Diabetes Endocrinol. 2015;3:286–95. https://doi.org/10.1016/ S2213-8587(14)70225-6.
- Ramalingaswami V, Subramanian TA, Deo MG. The aetiology of Himalayan endemic goitre. Lancet. 1961;277:791–4.
- Knudsen N, Laurberg P, Perrild H, Bulow I, Ovesen L, Jorgensen T. Risk factors for goiter and thyroid nodules. Thyroid. 2002;12:879–88. https://doi. org/10.1089/105072502761016502.
- Mukhopadhyay S, Ghosh D, Chatterjee A, Sinha S, Tripathy S, Chandra AK. Evaluation of possible goitrogenic and anti-thyroidal effect of nitrate, a potential environmental pollutant. Indian J Physiol Pharmacol. 2005;49:284–8.
- Lundberg JO, Weitzberg E. Nitric oxide signaling in health and disease. Cell. 2022;185:2853–78. https://doi.org/10.1016/j.cell.2022.06.010.
- Erzurum SC, Ghosh S, Janocha AJ, Xu W, Bauer S, Bryan NS, Tejero J, Hemann C, Hille R, Stuehr DJ, et al. Higher blood flow and circulating NO products offset high-altitude hypoxia among Tibetans. Proc Natl Acad Sci U S A. 2007;104:17593–8. https://doi.org/10.1073/pnas.0707462104.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

